

SECTION VI: DISCUSSION OF CONCLUSIONS AND IMPLICATIONS

Section I. Introduction and Literature Review

The information provided in Section I provides a basis for understanding the importance of mast in oak dominated forests in terms of potential impacts on wildlife population dynamics and oak reproduction. The data are more extensive than is required in the mast model as it is currently formulated; however, the algorithm may be altered to accommodate a higher degree of detail or applied to different ecosystems where the current formulation is not appropriate.

Section II. Probability and Quantity of Acorn Production: Parameters for the Landis Add-on Module

Section II presents a synthesis of available information on factors relevant to this mast production model in the form of developing and justifying parameters and applying these relationships in the program code of the mast model. The requirements and limitations of the landscape model LANDIS are also summarized. In general, data from local inventories are used to stochastically estimate the diameter, density, and species-group distribution of oak trees in canopy positions on landscape elements (cells) in consideration of differences among local land types. Variability among trees in inherent acorn production is factored and an estimate of an area-based mast production index is generated.

One relationship that is modeled, a prediction of tree size from the age of cell dominants is probably a weakness of the model. While the predictability of the relationship is improved over normal situations by being able to limit analysis of

inventory data to only upper canopy trees in dominant and co-dominant positions (since these trees produce the vast majority of acorns), the relationship is still tenuous. For example, while one might expect a large number of small, suppressed trees in a stand where the age of the dominants is relatively old, (*e.g.* 160 years), one might be less willing to accept the presence of large trees on young stands (*e.g.* 20 years). Due to the high degree of variability associated with the age-diameter relationship, this sort of situation is not rare in the mast model output. Therefore, either a better description of the relationship or a more subjective or idealized application of variability may be warranted.

The formulation of the model is specific to forests in southern Missouri. The algorithm may be used in other ecosystem types; however, it is probable that the parameters would need to be altered and some information provided in the LANDIS output that was ignored in my formulation might be re-considered for other situations. For example, I did not utilize the species presence-absence information since an analysis of the local data revealed that oaks were distributed uniformly and predictably though out the forests of southern Missouri. However, LANDIS, as parameterized for other ecosystems, factors many more species (He *et al.* 1999) and since more species are considered, the presence or absence of oaks may be more pertinent.

Another improvement of the mast model algorithm would be to link the estimations of tree-level data among iterations of the LANDIS output. Currently, the algorithm stochastically assigns tree density, diameter, species distribution, and inherent mast production characteristics independently among iterations. Thus, a cell may change in character between the 20th and 30th year iterations. To some degree the algorithm compensates for this effect since the stand-level mast index is estimated from the average

of many cells (a 30 ha stand is comprised of 333 cells). Setting the initial “seed” for the random number generator explicitly before running simulations ensures the same random numbers are assigned repeatedly. Thus, comparison among simulated treatments is not affected by stochastic variation among cells.

The relationship between tree size and mast production by species was scaled in a manner that produced a value that is roughly equivalent to a mass-per-unit-area value (kg acorns ha⁻¹). By using these two values interchangeably, I hope to convey the impression that the value is intended to be an index but may have a more tangible interpretation. Table VI.1 presents a summary from three sources of data for converting from the mass of acorns to the number of acorns by some species found in Missouri. If a modeler is interested in the number of acorns produced, the wide range in the relative weight of acorns (*e.g.* 0.4 to 3.1 g, air-dried) among species suggests that in the model, rather than distribution of trees among sub-genera, perhaps canopy trees should be distributed among species.

Section III. Implementation and Behavior of the Spatial Component of the Masting Model

The coding of the model seems to simulate the desired relationships in the model algorithm. The tree density (TPC) and species group assignment routines in general assign stand characteristics typical of southern Missouri. The diameter distributions are not as well defined and estimation may need to be improved (see above). The results of ultimate interest (the mast indices) followed the expected pattern with respect to age and species groups. That is, as stands grew older, trees of the white oak group became more important contributors to the mast crop.

Table VI.1 Some conversion factors for estimating the number and mass of fresh and dry acorns per unit area (Adapted from Waller 1979, Downs 1944, and the MOFEP hard mast plot data).

	Interval between large seed crops [years] (Waller)	Acorns per kg (Waller)	Acorn Mass [g] (Waller)	Acorn Mass [g] Fresh (Downs)	MOFEP		Estimated Conversion factor fresh to dry
					Air Dried Mass 93,95,97,99 [g acorn ⁻¹]	Acorns kg ⁻¹	
<i>Quercus alba</i>	7	264	3.8	2.7	1.1	891	0.41
<i>Q. coccinea</i>	4	517	1.9	2.3	1.4	690	0.62
<i>Q. falcata</i>	1.5	1188	0.8		0.4		
<i>Q. palustris</i>	1.5	902	1.1		0.6		
<i>Q. prinus</i>	2.5	220	4.5	6.6	2.3		
<i>Q. rubra</i>	4	275	3.6	5.1	3.1	325	0.61
<i>Q. velutina</i>	2.5	539	1.9	1.9	1.1	927	0.57
<i>Q. stellata</i>					0.4	2816	

Sensitivity analysis revealed that in terms of predicting cell mast index estimates, the parameters for the distribution of canopy trees among oak sub-genera were the most sensitive to change. The diameter-age relationship and the TPC estimation were not nearly as sensitive to change. This was perhaps due to the application of a relatively small amount of variation to the species-distribution estimate versus the large amount of variation factored in predicting tree size and stand-cell density.

Running the model on the 3,216 ha test landscape provided data for generating maps of mast production. A more parsimonious re-formulation of this first version of the mast model may improve performance in terms of computer time and memory usage. In many cases, temporary variables can be created and immediately discarded rather than retained in active memory and later discarded. Another approach for improving model efficiency, would be to parameterize the model for a situation and run the model on only a sub-sample of the entire landscape and estimate the mean and variation in the cell-level mast production index (CMPI) as a function of the age of dominants and land type. A separate and much simpler model could then apply this relationships ($CMPI = f(\text{age and landtype})$) stochastically across the entire landscape. Once summarized at the stand level, the results of the simpler model should resemble the results of the complete model.

Section IV. Implementation of Annual Variability in the Masting Model

The model adding annual variability to the spatial model seemed to produce results similar to those seen in nature. The model could be improved by having more explicit information regarding the probabilities of growing season factors that affect mast production (*e.g.* hail, wind, insect predation, drought etc). Also, for this model the effects

of microclimate are only hypothesized. I have subjectively estimated the probability of mast-limiting events and subjectively assigned the degree of influence these events have on different land types. This is clearly an opportunity for further research. The recently evolving availability of precociously fruiting oak trees that are small enough to enclose in greenhouses and growth chambers provides one such opportunity.

To look at how global climate change might affect mast production, I reviewed some internet-based sources of information and ran the mast model on one scenario. There is no general consensus regarding how the climate in the mid-west U.S. will be affected, particularly regarding relative humidity. However, one common prediction is that while, in general, the average temperatures will rise, the weather patterns will become more erratic and relative humidity will rise (Watson et al. 1997; EPA 2000; Green 2000, WMO 2001). For my simulation, I interpret this to mean that, i) since flowering is triggered by cumulative heating in the spring, flowers may mature earlier but since the climate is more erratic, there is a higher probability of freezing damage, and ii) higher relative humidity will increase the likelihood of poor flowering conditions. Figure VI.1 depicts the amount of the change of influence on freezing probability and relative humidity effect that I factored in this simulation. Basically, the probability of freezing was doubled and the influence of relative humidity altered to a lesser extent. The simulation was run on the output of the spatial mast model for the even-aged management 90th year iteration with and without the climate change using the same initial random number seed. The results suggest a relatively small impact of climate change (given this scenario) on total mast production (Figure VI.2). There were some years (2,8,9,10) of the simulation that under the base climate parameters produced a small




Figure VI.1 Change in the scale factors under the climate change scenario. A random number is generated and the influence of the factor (scale multiplier) is derived from the relationships shown.




Figure VI.2 The change in mast production under a climate change scenario that increased the frequency of freezing events during flowering stages and increased probability of high relative humidity.

amount of mast ($<50 \text{ kg ha}^{-1}$) while during those same years little or no mast production occurred. Most high mast production years were not affected at all.

Section V. Allocation of Resources: The Trade-off Between Vegetative Growth and Acorn Production.

The results of the study of resource allocation to mast production suggest that, although vague, there is reason to suspect that there is a measurable trade-off in resources between acorn production and diameter and following year shoot-length growth. As mentioned above, there may be an opportunity for studying this possible relationship using precociously flowering and fruiting oak trees. Early fruiting is an inherited trait and may also be an indicator of overall tree vigor.

CONCLUSION

The masting model algorithm provides a framework for adding tree-level detail to a landscape model (LANDIS). A mechanistic approach was adopted to assign characteristics to cells that might affect mast production. These characteristics were then used to estimate mast production. While the parameters and formulation would change, this general approach can be used to address other research questions mechanistically.

Literature Cited

- Auchmoody, L.R., H.C. Smith, and R.S. Walters. 1993. Acorn production in northern red oak stands in northwestern Pennsylvania. Res. Pap. NE-680. USDA Forest Service, Northeast Forest Experiment Station. 5 p.
- Bartelink, H.H. 1998. A model of dry matter partitioning in trees. *Tree Physiology*. 18: 91-101.
- Brookshire, B.L. and S.R. Shifley. 1997. Proceedings of the Missouri Ozark Forest Ecosystem Project: Symposium. Gen. Tech. Rep. NC-193. USDA Forest Service, North Central Research Station. St. Paul, MN. 378 p.
- Cecich, R. A. 1992. Flowering and oak regeneration: IN: Loftis, D.L. and C.E. McGee (eds), *Oak Regeneration: Serious Problems, Practical Recommendations*. Sept 8-10, 1992. USDA. Forest Service. Gen. Tech. Rep. SE-84. Southeastern Forest Experiment Station, Asheville, N.C.
- Cecich, R.A. 1994. The reproductive biology of *Quercus*, with an emphasis on *Q. rubra*. In: *Biology and silviculture of northern red oak in the north central region: A synopsis*. J.G. Isebrands and R.E. Dickson (eds) USDA Forest Service Gen. Tech. Rep. NC-173. 68 p.
- Cecich, R.A. and N.H. Sullivan. 1999. Influence of weather at time of pollination on acorn production of *Quercus alba* and *Quercus velutina*. *Can J. For. Res.* 29: 1817-1823.
- Dickson, J.G. 1990. Oak and flowering dogwood fruit production for eastern wild turkeys. IN: W.M. Healy, G.B. Healy, eds. Proc. 6th National Wild Turkey Symposium: National Wild Turkey Federation: 90-95.
- Dickson, R.E. and P.T Tomlinson. 1996. Oak growth, development and carbon metabolism in response to water stress. *Ann. Sci. For.* 53: 181-196.
- Downs, A.A. 1944. Estimating acorn crops for wildlife in the southern Appalachians. *J. of Wildlife Management*. 8(4): 339-340.
- Downs, A.A. 1949. Trees and food from acorns. *USDA Yearbook of Agriculture*, 1949. 571-573.
- Downs, A.A., and W.E. McQuilken. 1944. Seed production of southern Appalachian oaks. *J. of Forestry*. 42(12): 913-920.
- Drake, W.E. 1991. Evaluation of an approach to improve acorn production during thinning. IN: Proceedings of the 8th Central Hardwood Conference. L.H. McCormick and K.W. Gottschalk (eds). USDA Forest Service. Gen Tech. Rep. NE-148. Radnor, PA 605 p.
- EPA 2000. Global Warming Site: Climate [http:// ww.epa.gov/globalwarming/climate/index.html](http://ww.epa.gov/globalwarming/climate/index.html)
- Farmer, R.E. 1981. Variation in seed yield of white oak. *Forest Science*. 27(2):377-380.

- Feret, P.P. , R.E. Kreh, S.A. Merckle, and R.G. Oderwald. 1982. Flower abundance, premature acorn abscission, and acorn production in *Quercus alba* L. *Botanical Gazette*. 143(2): 216-218.
- Goodrum, P.D., V.H. Reid, and C.E. Boyd. 1971. Acorn yields, characteristics, and management criteria of oaks for wildlife. *J. of Wildlife Management*. 35(3): 520-532.
- Green. K. 2000. A Plain English Guide to Climate Change.
<http://www.rpi.org/environment/peg3.html>
- Greenberg, C.H. 2000. Individual variation in acorn production by five species of southern Appalachian oaks. *Forest Ecology and Management*. 132: 199-210.
- Gustafson, E. J., S.R. Shifley, D.J. Mladenoff, K.K. Nimerfro, and H.S. He. 2000. Spatial simulation of forest succession and timber harvesting using LANDIS. *Can. J. For. Res.* 30: 32-43.
- Gysel, L.W. 1957. Acorn production on good, medium, and poor oak sites in southern Michigan. *J. Forestry*. 55: 570-574.
- Hannon, S.J., R.L. Mumme, W.D. Koenig, S. Spon, and F.A. Pitelka. 1987. Poor acorn crop, dominance, and decline in numbers of acorn woodpeckers. *J. of Animal Ecology*. 56: 197-207.
- Hansen, M.H.,T. Frieswyk, J.F. Glover, and J.F. Kelly. 1992. The Eastwide Forest Inventory Data Base: Users Manual. Gen. Tech. Rep. NC-151. USDA Forest Service. St. Paul, MN. 48 p. Internet Address: (<http://www.srsfia.usfs.msstate.edu/scripts/ew.htm>).
- Harlow, R.F. and R.L. Eikum. 1963. The effect of stand density on the acorn production of turkey oaks. Proc. 17th Ann. Conf. Southeast Assoc. Game and Fish Commisioners. 126-133.
- Harlow, R.F., J.B. Whelan, H.S. Crawford, and J.E. Skeen. 1975. Deer food during years of oak mast abundance and scarcity. *J. of Wildlife Management*. 39(2): 330-336.
- Healy, W.M. 1997. Thinning New England oak stands to enhance acorn production. *North. J. Applied Forestry*. 14(3): 152-156.
- Healy, W.M., A.N. Lewis, and E.F. Boose. 1999. Variation of red oak acorn production. *Forest Ecology and Management*. 116: 1-11.
- Hong, HS, D.J. Mladenoff, T.R. Crow. 1999. Linking an ecosystem model and a landscape model to study forest species response to climate warming. *Ecological Modelling* 114. 213-233.
- Hong, H.S., and D.J. Mladenoff. 1999. Spatially explicit and stochastic simulation of forest-landscape fire disturbance and succession. *Ecology*. 80(1): 81-99.
- Innouye D.W., D.E. Gill, M.R. Dudash, and C.B. Fenster. 1994. A model and lexicon for pollen fate. *American J. of Botany*. 81(12):1517-1530.
- Johnson, F.L. 1975. White oak acorn production in the upland streamside forest of central Illinois. Univ. Illinois. Agric. Exp. Stn. For Res. Rep. 75-3.

- Johnson, P.J. 1990. *Quercus coccinea* IN: Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.
- Koenig, W.D. W.J. Carmen, M.T. Stanback, and R.L. Mumme. 1991. Determinants of acorn productivity among five species of oaks in central coastal California. USDA For. Serv. Gen. Tech. Rep. PSW-126 pp 136-142.
- Koenig, W.D., R.L. Mumme, W.J. Carmen, and M.T. Stanback. 1994. Acorn production by oaks in central coastal California: Variation within and among years. *Ecology*. 75(1):99-109.
- Krajicek, J.E., K.A. Brinkman, and S.F. Gingrich. 1961. Crown competition- A measure of density. *Forest Science* 7(1): 35-42.
- LaLonde, R.G., and B.D. Roitberg. 1992. On the evolution of masting behavior in trees: predation or weather. *American Naturalist*. 139(6): 1293-1304.
- Liljegren et al. 1999. Interactions among a APETALA1, LEAFY, TERMINAL FLOWER1 specify meristem fate. *The Plant Cell*. 11: 1007-1018.
- Loewenstein, E.F., P.S. Johnson, and H.E. Garrett. 2000. Age and diameter structure of a managed uneven-aged oak forest. *Canadian J. of Forest Research*. 30(7):1060-1070.
- Makela, A. 1997. A carbon balance model of growth and self-pruning in trees based on structural relationships. *Forest Science*. 43(1): 7 – 23.
- Malstrom, H.L, and J.L. McMeans. 1982. Shoot length and previous fruiting affect subsequent growth and nut production of 'Moneymaker' Pecan. *HortScience*. 17(6): 970-972.
- McQuilkin, R.A. 1990. *Quercus palustris* : IN Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.
- McQuilkin, R.A. 1990. *Quercus prinus* : Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.
- McShea, W.J. and G. Schwede. 1993. Variable acorn crops: Responses of white-tailed deer and other mast consumers. *J. Mammology*, 74(4): 999-1006.
- Minckler, L.S. and D. Janes. 1965. Pin oak acorn production on normal and flooded areas. University of Missouri. Agric. Exp. Stn. Research Bull. 898.
- Minckler, L.S. and R.F. McDermott. 1960. Pin oak acorn production and regeneration as affected by stand density, structure and flooding. University of Missouri. Agric. Exp. Stn. Research Bull. 750.
- Mizukami and Ma. 1997. Determination of *Arabidopsis* floral meristem identity by Agamous. *The Plant Cell*. 9: 393-408.

Mladenoff, D.J., G.E. Host, J. Boeder, and T.R. Crow. 1996. LANDIS: a spatial model of forest landscape disturbance. Succession, and management. In: M.F. Goodchild et al. (eds). GIS and Environmental Modeling: Progress and Research Issues. GIS World Books, Fort Collins, CO. P. 175 - 180.

Mladenoff, D.J. and H.S. He. 1999. Design and behavior of LANDIS, an object oriented model of forest landscape disturbance and succession. In: Mladenoff, D.J., W.L. Baker (eds). Advances in Spatial Modeling of Forest Landscape Change: Approaches and Applications. Cambridge, UK. Cambridge University Press. p 125-162.

Myers, S.A. 1978. *Insect impact on acorn production in Missouri upland forests*. Thesis (Ph.D.) University of Missouri, CO. 245 p.

Nixon, C.M., and M.W. McClain. 1969. Squirrel population decline following late spring frost. *J. of Wildlife Management*. 33(2): 353-357.

Norton, D.A. and D. Kelly. 1988. Mast seeding over 33 years by *Darydium cupressinum* Lamb. (rimu)(*Podocarpaceae*) in New Zealand: the importance of economies of scale. *Functional Ecology*. 2: 399-408.

Ostfeld, R.S. C.G. Jones and J.O. Wolff. 1996. Of mice and mast: Ecological connections in eastern deciduous forests. *BioScience*. 46(5):323-330.

Post, L.S. 1998. Seed management in Tennessee: Development of seed zones for Tennessee and distribution and protection of northern red oak (*Quercus rubra* L.) acorns. Knoxville, TN: University of Tennessee. 177 p. Thesis.

Potter, B.E. and T.W. Cate. 1999. A climatology of late-spring freezes in the northeastern United States. USDA Forest Service GTR NC-204. St. Paul. MN. 35 p.

Reekie, E.G. and F.A. Bazzaz. 1987. Reproductive effort in plants. 1. Carbon allocation to reproduction. *American Naturalist*. 129(6): 876-896.

Rogers, R.R. 1990. *Quercus alba*. IN: Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.

Sander, I.L. 1990. *Quercus rubra*: IN. Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.

Sander, I.L. 1990. *Quercus velutina*: IN. Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.

Seki, T. 1994. Dependency of cone production on tree dimensions in *Abies mariesii*. *Canadian J. Botany* 72: 1713-1719.

Sharp and Chisman 1961. Flowering and fruiting in the white oaks. I. Staminate flowering through pollen dispersal. *Ecology*. 42: 365-372.

- Sharp, W.M. and V.G. Sprague. 1967. Flowering and fruiting in the white oaks. Pistillate flowering, acorn development, weather, and yields. *Ecology*. 48(2):243-251.
- Shelburne, D.R. 1983. Patterns in the production, development, and maturation of acorns in *Quercus alba* (Fagaceae). MS Thesis. University of Missouri-St. Louis. 80 p.
- Shifley, S.R. and B.L. Brookshire. 2000. Missouri Ozark Forest Ecosystem Project: site history, soils, landforms, woody and herbaceous vegetation, down wood, and inventory methods for the landscape experiment. Gen. Tech. Rep. NC-208. USDA Forest Service, North Central Research Station. St. Paul, MN. 314 p.
- Shifley, S.R., F.R. Thompson, D.R. Larsen, and W.D. Dijak. 2000. Modeling forest landscape change in the Missouri Ozarks under alternative management practices. *Computers and electronics in Agriculture*. 27(2000) 7-24.
- Silvertown, J.W. 1980. The evolutionary ecology of mast seeding in trees. *Biol. J. Linn. Soc.* 14:235-250.
- Sork, V.L., J. Bramble, and O. Sexton. 1993. Ecology of mast-fruited in three species of North American deciduous oaks. *Ecology*. 74(2): 528-541.
- Sprugel, D.G. T.M. Hinckley, and W. Schaap. 1991. The theory and practice of branch autonomy. *Annual Review of Ecological Systematics*. 22: 309-334.
- StataCorp. 1999. Stata Statistical Software: Release 6.0. College Station, TX: Stata Corporation.
- Stransky, J.J. 1990. *Quercus stellata*: IN. Silvics of North America. Vol 2. USDA Forest Service AG Handbook 654.
- Uhlig, H.G. and H.L. Wilson. 1952. A method of evaluating an annual mast index. *J. of Wildlife Management*. 16(3): 338-343.
- Waller, D.M. 1979. Models of mast fruiting in trees. *J. of Theoretical Biology*. 80: 223-232.
- Williamson, M.J. 1966. Premature Abscissions and white oak acorn crops. *Forest Science*. 12(1): 19-21.
- Wilson, P.R., B.J. Karl, R.J. Toft, J.R. Beggs, and R.H. Taylor. 1998. The role of introduced predators and competitors in the decline of Kaka (*Nestor meridionalis*) Populations in New Zealand. *Biological Conservation*. 83 (2): 175-185.
- Wolff, J.O. 1996. Population fluctuations of mast-eating rodents are correlated with production of acorns. *J. Mammology*. 77(3): 850-856.
- Wolgast, L.J. and B.B. Stout. 1977. The effects of relative humidity at the time of flowering on fruit set in bear oak (*Quercus ilicifolia*) mer. *J. Botany*. 64 (2): 159-160.

- Yanofsky. 1995. Floral Meristems to floral organs: Genes controlling early events in *Arabidopsis* flower development. *Ann. Rev. Physiol. Plant. Mol. Biol.* 46: 167-188.
- Valentine, H.T., T.G. Gregoire, H.E. Burkhart, and D.Y. Hollinger. 1997. A stand-level model of carbon allocation and growth, calibrated for loblolly pine. *Canadian J. Forest Research.* 27: 817-830.
- VanGilder, L.D. 1997. Acorn production on the Missouri Forest Ecosystem project study sites: Pre-treatment data. IN: Brookshire, B.L. and S.R. Shifley (eds.) *Proceedings of the Missouri Forest Ecosystem Project Symposium: an experimental approach to landscape research*: USDA Forest Service. GTR NC-193. St. Paul, MN 378 p.
- Vogt, K. 1991. Carbon budgets of temperate forest ecosystems. *Tree Physiology.* 9: 69-86.
- Waring, R.H. and W.H. Schlesinger. 1985. *Forest Ecosystems: Concepts and Management*. Academic Press, Inc. New York. 340 p.
- Watson, R.T. , M.C. Zinyowera, R.H. Moss, and D.J. Dokken. 1997. The Regional Impacts of Climate Change: an assessment of vulnerability. [http:// www.ipcc.ch/pub/sr97.htm](http://www.ipcc.ch/pub/sr97.htm)
- WMO 2001 World Meteorological Organization.
<http://www.wmo.ch/web/gcos/terre/variable/relhum.htm> Geneva, Switzerland

VITA

Neal Sullivan was born March 7, 1960 in Peabody, Massachusetts. His first exposure to natural resource issues in an academic setting was at Essex Agricultural and Technical Institute in Hathorne, Massachusetts where he attended high school. After graduating, he entered the military, was stationed in Germany and after three years, returned to the U.S. to continue his education. He received the following degrees: an A.A.S. in Natural Resources Management from Essex Agricultural and Technical Institute at Hathorne, Massachusetts (1985) ; a B.S.-Forestry and an MSc. in Natural Resources from the University of New Hampshire at Durham (1989 and 1993, respectively): and a Ph.D. in Forestry from the University of Missouri at Columbia (2001). He is married to Michelle Boone formerly of Goose Creek, South Carolina and is presently a Post-Doctoral Research Fellow with the Department of Forestry at the University of Missouri, Columbia.